

Geometric Models for Secondary Structures in Proteins

Magdalena Toda

*Texas Tech University, Department of Mathematics and Statistics, 2500
Broadway Ave, Lubbock, Texas, 79409-1042, USA*
E-mail:magda.toda@ttu.edu

ABSTRACT

This reports represents a study of beta barrels as a secondary structure in proteins, using methods from differential geometry and variational calculus, namely Dirichlet and Willmore-type energies. We review some historical models of beta sheets and barrels based on best-fitting hyperboloids and catenoids, respectively, and explain why those models are outdated. We provide an elastic membrane model for these structures, via a Willmore type energy that is similar to the Helfrich energy for lipid bilayers.

REFERENCES

- [1] Vogel T., *Stability of a Liquid Drop Trapped Between Two Parallel Planes*, SIAM Journal on Applied Mathematics **47 - 3** (1987) 516 - 525.
- [2] Koh E., Kim T., *Minimal Surface as a Model of Beta-Sheets*, Proteins: Structure, Function and Bioinformatics **61** (2005) 559 - 569.
- [3] Novotny J., *Twisted Hyperboloid (Strophoid) as a Model of Beta-Barrels in Proteins*, Molecular Biology **177** (1984) 567 - 573.
- [4] Lasters I. and all, *Structural Principles of Parallel Beta-Barrels in Proteins*, Proc. National Academy of Science USA **85** (1988) 3338 - 3342.
- [5] Helfrich W., *Elastic Properties of Lipid Bilayers: Theory and Possible Experiments*, Zeitschrift für Naturforschung, Teil C: Biochemie, Biophysik, Biologie, Virologie **28-11** (1973) 693.
- [6] Hiroyoshi N., Masahiro O., Ou-Yang Z., *Counterexample to Some Shape Equations for Axisymmetric Vesicles*, Physical Review E **48** (1993) 2304 - 2307.