Fifteenth International Conference on Geometry, Integrability and Quantization June 7–12, 2013, Varna, Bulgaria Ivaïlo M. Mladenov, Andrei Ludu and Akira Yoshioka, Editors Avangard Prima, Sofia 2014, pp 218–241 doi: 10.7546/oic-15-2014-218-241



ESSENTIAL NONLINEARITY IN FIELD THEORY AND CONTINUUM MECHANICS. SECOND- AND FIRST-ORDER GENERALLY-COVARIANT MODELS

EWA ELIZA ROŻKO and JAN JERZY SŁAWIANOWSKI

Institute of Fundamental Technological Research, Polish Academy of Sciences 5^{B} , Pawińskiego Str., 02-106 Warsaw, POLAND

Abstract. Discussed is the problem of the mutual relationship of differentially first-order and second-order field theories and quantum-mechanical concepts. We show that unlike the real history of physics, the theories with algebraically second-order Lagrangians are primary, and in any case more adequate. It is shown that in principle, the primary Schrödinger idea about Lagrangians which are quadratic in derivatives, and leading to second-order differential equations, is not only acceptable, but just it opens some new perspective in field theory. This has to do with using the Lorentz-conformal or rather its universal covering $\mathrm{SU}(2,2)$ as a gauge group. This has also some influence on the theory of defects in continua.

CONTENTS

1.	Summary	219
2.	Heuristics of the $\mathrm{U}(1)$ Gauge Group	219
3.	SU(2, 2) as a Gauge Group	221
4.	Second-Order $\mathrm{SU}(2,2)$ Gauge Theory	229
5.	The $\mathrm{SL}(2,C)$ - Correspondence in the Second-Order $\mathrm{SU}(2,2)$ Theory	232
6.	First-Order $\mathrm{SU}(2,2)$ Field Theory	
Ac	knowledgements	240
References		240