

ON THE DECOMPOSITION
OF THE OSCILLATOR REPRESENTATION

ALEKSANDER STRASBURGER
*University of Warsaw, Faculty of Physics,
Department of Mathematical Methods of Physics
Hoża 74, 00-682 Warszawa, Poland*

Following Howe [7] we shall call the oscillator representation in \mathbf{R}^d the representation of the Lie algebra $f\mathfrak{U}(\epsilon, \mathbf{R})$ on the Schwartz space $S(\mathbf{R}^d)$ or the corresponding (i.e. exponentiated) unitary representation on $L^2(\mathbf{R}^d)$ of the double cover $\widetilde{SL}(2, \mathbf{R})$ of $SL(2, \mathbf{R})$ obtained in the following way. Let

$$h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad e^+ = \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad e^- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

be a standard basis of the Lie algebra $sl(2, \mathbf{R})$ satisfying the commutation relations

$$[h, e^\pm] = \pm 2e^\pm, \quad [e^+, e^-] = h.$$

We let $\Delta = \sum_{j=1}^d \frac{\partial^2}{\partial x_j^2}$ stand for the Laplace operator and we shall use the familiar notation $r^2 = r^2(x) = x \cdot x$ for the function on \mathbf{R}^d equal to the square of the radius (for the standard euclidean metric). If D is a differential operator (e.g. the Laplacian) and f a smooth function, we shall write $D \circ f$ for the composition of D with the operator of multiplication by f . The oscillator representation ω^d of the Lie algebra $sl(2, \mathbf{R})$ is defined by setting

$$\begin{aligned} \omega^d(h) &= \sum_{j=1}^d x_j \frac{\partial}{\partial x_j} + \frac{d}{2}, \\ \omega^d(e^+) &= \frac{i}{2} \sum_{j=1}^d x_j^2 = \frac{i}{2} r^2, \\ \omega^d(e^-) &= \frac{i}{2} \sum_{j=1}^d \frac{\partial^2}{\partial x_j^2} = \frac{i}{2} \Delta \end{aligned} \tag{1}$$

and extending by linearity to the whole of $sl(2, \mathbf{R})$. The fact that this representation of the Lie algebra $sl(2, \mathbf{R})$ is a derived representation of a unitary representation of the double cover $\widetilde{SL}(2, \mathbf{R})$ of $SL(2, \mathbf{R})$ is a part of a theorem of Shale and A. Weil asserting the existence of the metaplectic representation, cf. e.g. [6] or [3].

Many properties of the oscillator representation are easily accessed through the use of another basis satisfying the standard commutation relations, namely the one given by the matrices

$$k = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \quad n^+ = \frac{1}{2} \begin{pmatrix} 1 & i \\ i & -1 \end{pmatrix}, \quad n^- = \frac{1}{2} \begin{pmatrix} 1 & -i \\ -i & -1 \end{pmatrix},$$

which satisfy

$$[k, n^\pm] = \pm 2n^\pm, \quad [n^+, n^-] = k.$$